

CONTINUED FRACTIONS AND GENERALIZED PATTERNS

TOUFIK MANSOUR

LABRI, Université Bordeaux I,
351 cours de la Libération, 33405 Talence Cedex
toufik@labri.fr

ABSTRACT

In [BS] Babson and Steingrímsson introduced generalized permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation.

Let $f_{\tau;r}(n)$ be the number of 1-3-2-avoiding permutations on n letters that contain exactly r occurrences of τ , where τ a generalized pattern on k letters. Let $F_{\tau;r}(x)$ and $F_{\tau}(x, y)$ be the generating functions defined by $F_{\tau;r}(x) = \sum_{n \geq 0} f_{\tau;r}(n)x^n$ and $F_{\tau}(x, y) = \sum_{r \geq 0} F_{\tau;r}(x)y^r$. We find an explicit expression for $F_{\tau}(x, y)$ in the form of a continued fraction for where τ given as a generalized pattern; $\tau = 12\text{-}3\text{-}\dots\text{-}k$, $\tau = 21\text{-}3\text{-}\dots\text{-}k$, $\tau = 123\dots k$, or $\tau = k\dots 321$. In particular, we find $F_{\tau}(x, y)$ for any τ generalized pattern of length 3. This allows us to express $F_{\tau;r}(x)$ via Chebyshev polynomials of the second kind, and continued fractions.

1. INTRODUCTION

Let $[p] = \{1, \dots, p\}$ denote a totally ordered alphabet on p letters, and let $\pi = (\pi_1, \dots, \pi_m) \in [p_1]^m$, $\beta = (\beta_1, \dots, \beta_m) \in [p_2]^m$. We say that π is *order-isomorphic* to β if for all $1 \leq i < j \leq m$ one has $\pi_i < \pi_j$ if and only if $\beta_i < \beta_j$. For two permutations $\pi \in S_n$ and $\tau \in S_k$, an *occurrence* of τ in π is a subsequence $1 \leq i_1 < i_2 < \dots < i_k \leq n$ such that $(\pi_{i_1}, \dots, \pi_{i_k})$ is order-isomorphic to τ ; in such a context τ is usually called the *pattern* (classical pattern). We say that π *avoids* τ , or is τ -*avoiding*, if there is no occurrence of τ in π . More generally, we say π *containing* τ exactly r times, if there exists r different occurrences of τ in π .

The set of all τ -avoiding permutations of all possible sizes including the empty permutation is denoted $\mathcal{S}(\tau)$. Pattern avoidance proved to be a useful language in a variety of seemingly unrelated problems, from stack sorting [Kn] to singularities of Schubert varieties [LS]. A complete study of pattern avoidance for the case $\tau \in S_3$ is carried out in [SS].

On the other hand, in [BS] introduced generalised permutation patterns that allow the requirement that two adjacent letters in a pattern must be adjacent in the permutation. The idea of [BS] introducing these patterns was study of Mahonian statistics.

We write a classical pattern with dashes between any two adjacent letters of the pattern, say 1324, as 1-3-2-4, and if we write, say 24-1-3, then we mean that if this pattern occurs in permutation π , then the letters in the permutation π that correspond to 2 and 4 are adjacent. For example, the permutation $\pi = 35421$ has

only two occurrences of the pattern 23-1, namely the subsequences 352 and 351, whereas π has four occurrences of the pattern 2-3-1, namely the subsequences 352, 351, 342 and 341.

In [C] presented a complete solution for the number of permutations avoiding any pattern of length three with exactly one adjacent pair of letters. In [CM] presented a complete solution for the number of permutations avoiding any two patterns of length three with exactly one adjacent pair of letters. In [Ki] presented almost results avoiding two or more 3-patterns without internal dashes, that is, where the pattern corresponds to a contiguous subword in a permutation. Besides, [EN] presented the following generating functions regarding the distribution of the number of occurrences of any generalized pattern of length 3:

$$\begin{aligned} \sum_{\pi \in S} y^{(123)\pi} \frac{x^{|\pi|}}{|\pi|!} &= \frac{2f(y)e^{\frac{1}{2}(f(y)-y+1)x}}{f(y)+y+1+(f(y)-y-1)e^{f(y)x}}, \\ \sum_{\pi \in S} y^{(213)\pi} \frac{x^{|\pi|}}{|\pi|!} &= \frac{1}{1 - \int_0^x e^{(y-1)t^2/2} dt}, \end{aligned}$$

where $(\tau)\pi$ is the number of occurrences of τ in π , $f(y) = \sqrt{(y-1)(y+3)}$.

The purpose of this paper is to point out an analog of a theorem [RWZ], and some interesting consequences of this analog. Generalization of this theorem have already been given in [Kr, MV, JR]. In the present note we study the generating function for the number 1-3-2-avoiding permutations in S_n that contain a prescribed number of generalised pattern τ . The study of the obtained continued fraction allows us to recover and to present analog of the results of [CW, Kr, MV, JR] that relates the number of 1-3-2-avoiding permutations that contain no 12-3-...- k (or 21-3-...- k) patterns to Chebyshev polynomials of the second kind.

Let $f_{\tau;r}(n)$ stand for the number of 1-3-2-avoiding permutations in S_n that contain exactly r occurrences of τ . We denote by $F_{\tau;r}(x)$ and $F_{\tau}(x, y)$ the generating function of the sequence $\{f_{\tau;r}(n)\}_{n \geq 0}$ and $\{f_{\tau;r}(n)\}_{n,r \geq 0}$ respectively, that is,

$$F_{\tau;r}(x) = \sum_{n \geq 0} f_{\tau;r}(n)x^n, \quad F_{\tau}(x, y) = \sum_{r \geq 0} F_{\tau;r}(x)y^r.$$

The paper is organized as follows. The cases $\tau = 12-3-...-k$, $\tau = 21-3-...-k$, $\tau = 123...k$, and $\tau = k...321$ are treat in **section 2**. In **section 3**, we treat the cases $\tau = 123, 213, 231, 312$, and 321 , that is τ is 3-letters generalized pattern without dashes. In **section 4**, we treat the cases when τ is 3-letters generalized pattern with one dash. Finally, in **section 5**, we treat some of examples of restricted more than one generalized pattern of 3-letters.

2. FOUR GENERAL CASES

In this section, we study the following four cases $\tau = 12-3-...-k$, $\tau = 21-3-...-k$, $\tau = 12...k$, and $\tau = k...21$, by the following three subsections.

2.1 PATTERN 12-3-...- k

Our first result is a natural analog of the main theorems of [RWZ, MV, Kr].

Theorem 2.1. *The generating function $F_{12-3-\dots-k}(x, y)$ for $k \geq 2$ is given by the continued fraction*

$$\cfrac{1}{1 - x + xy^{d_1} - \cfrac{xy^{d_1}}{1 - x + xy^{d_2} - \cfrac{xy^{d_2}}{1 - x + xy^{d_3} - \cfrac{xy^{d_3}}{\dots}}}},$$

where $d_i = \binom{i-1}{k-2}$, and $\binom{a}{b}$ is assumed 0 whenever $a < b$ or $b < 0$.

Proof. Following [MV] we define $\eta_j(\pi)$, $j \geq 3$, as the number of occurrences of $12-3-\dots-j$ in π . Define $\eta_2(\pi)$ for any π , as the number of occurrences of 12 in π , $\eta_1(\pi)$ as the number letters of π , and $\eta_0(\pi) = 1$ for any π , which means that the empty pattern occurs exactly once in each permutation. The *weight* of a permutation π is a monomial in k independent variables $q_1, \dots, q_k$ defined by

$$w_k(\pi) = \prod_{j=1}^k q_j^{\eta_j(\pi)}.$$

The *total weight* is a polynomial

$$W_k(q_1, \dots, q_k) = \sum_{\pi \in \mathcal{S}(1-3-2)} w_k(\pi).$$

The following proposition is implied immediately by the definitions.

Proposition 2.2. $F_{12-3-\dots-k}(x, y) = W_k(x, 1, \dots, 1, y)$ for $k \geq 2$.

We now find a recurrence relation for the numbers $\eta_j(\pi)$. Let $\pi \in S_n$, so that $\pi = (\pi', n, \pi'')$.

Proposition 2.3. *For any nonempty $\pi \in \mathcal{S}(1-3-2)$*

$$\eta_j(\pi) = \eta_j(\pi') + \eta_j(\pi'') + \eta_{j-1}(\pi'),$$

where $j \neq 2$. Besides, if π' nonempty then

$$\eta_2(\pi) = \eta_2(\pi') + \eta_2(\pi'') + 1,$$

otherwise

$$\eta_2(\pi) = \eta_2(\pi'').$$

Proof. Let $l = \pi^{-1}(n)$. Since π avoids $1-3-2$, each number in π' is greater than any of the numbers in π'' . Therefore, π' is a $1-3-2$ -avoiding permutation of the numbers $\{n-l+1, n-l+2, \dots, n-1\}$, while π'' is a $1-3-2$ -avoiding permutation of the numbers $\{1, 2, \dots, n-l\}$. On the other hand, if π' is an arbitrary $1-3-2$ -avoiding permutation of the numbers $\{n-l+1, n-l+2, \dots, n-1\}$ and π'' is an arbitrary $1-3-2$ -avoiding permutation of the numbers $\{1, 2, \dots, n-l\}$, then $\pi = (\pi', n, \pi'')$ is $1-3-2$ -avoiding. Finally, if $(i_1, \dots, i_j)$ is an occurrence of $12-3-\dots-j$ in π then either $i_j < l$, and so it is also an occurrence of $12-3-\dots-j$ in π' , or $i_1 > l$, and so it is also an occurrence of $12-\dots-j$ in π'' , or $i_j = l$, and so $(i_1, \dots, i_{j-1})$ is an occurrence of $12-3-\dots-(j-1)$ in π' , where $j \neq 2$. For $j = 2$ the proposition is trivial. The result follows. $\square$

Now we are able to find the recurrence relation for the total weight W . Indeed, by Proposition 2.3,

$$\begin{aligned} W_k(q_1, \dots, q_k) &= 1 + \sum_{\emptyset \neq \pi \in \mathcal{S}(1-3-2)} \prod_{j=1}^k q_j^{\eta_j(\pi)} \\ &= 1 + \sum_{\emptyset \neq \pi' \in \mathcal{S}(1-3-2)} \sum_{\pi'' \in \mathcal{S}(1-3-2)} \prod_{j=1}^k q_j^{\eta_j(\pi'')} \cdot q_1^{\eta_1(\pi')+1} q_2 \cdot \\ &\quad \prod_{j=2}^{k-1} (q_j q_{j+1})^{\eta_j(\pi')} \cdot q_k^{\eta_k(\pi')} + \sum_{\pi'' \in \mathcal{S}(1-3-2)} q_1 \prod_{j=1}^k q_j^{\eta_j(\pi'')}. \end{aligned}$$

Hence

$$\begin{aligned} W_k(q_1, \dots, q_k) &= 1 + q_1 W_k(q_1, \dots, q_k) + \\ &\quad + q_1 q_2 W_k(q_1, \dots, q_k) (W_k(q_1, q_2 q_3, \dots, q_{k-1} q_k, q_k) - 1). \end{aligned} \quad (1)$$

Following [MV], for any $d \geq 0$ and $2 \leq m \leq k$ define

$$\mathbf{q}^{d,m} = \prod_{j=2}^k q_j^{\binom{d}{j-m}};$$

recall that $\binom{a}{b} = 0$ if $a < b$ or $b < 0$. The following proposition is implied immediately by the well-known properties of binomial coefficients.

Proposition 2.4. *For any $d \geq 0$ and $2 \leq m \leq k$*

$$\mathbf{q}^{d,m} \mathbf{q}^{d,m+1} = \mathbf{q}^{d+1,m}.$$

Observe now that $W_k(q_1, \dots, q_k) = W_k(q_1, \mathbf{q}^{0,2}, \dots, \mathbf{q}^{0,k})$ and that by (1) and Proposition 2.4

$$\begin{aligned} W_k(q_1, \mathbf{q}^{d,2}, \dots, \mathbf{q}^{d,k}) &= 1 + q_1 W_k(q_1, \mathbf{q}^{d,2}, \dots, \mathbf{q}^{d,k}) + \\ &\quad + q_1 q_2 W_k(q_1, \mathbf{q}^{d,2}, \dots, \mathbf{q}^{d,k}) (W_k(q_1, \mathbf{q}^{d+1,2}, \dots, \mathbf{q}^{d+1,k}) - 1). \end{aligned}$$

therefore

$$W_k(q_1, \dots, q_k) = \frac{1}{1 - q_1 + q_1 \mathbf{q}^{0,2} - \frac{q_1 \mathbf{q}^{0,2}}{1 - q_1 + q_1 \mathbf{q}^{1,2} - \frac{q_1 \mathbf{q}^{1,2}}{1 - q_1 + q_1 \mathbf{q}^{2,2} - \frac{q_1 \mathbf{q}^{2,2}}{\dots}}}}.$$

To obtain the continued fraction representation for $F(x, y; k)$ it is enough to use Proposition 2.2 and to observe that

$$q_1 \mathbf{q}^{d,2} \Big|_{q_1=x, q_2=\dots=q_{k-1}=1, q_k=y} = xy \binom{d}{k-2}.$$

□

Corollary 2.5.

$$F(x, y; 2) = \frac{1 - x + xy - \sqrt{(1-x)^2 - 2x(1+x)y + x^2 y^2}}{2xy},$$

in other words, for any $r \geq 1$

$$f_{12}(n) = \frac{r+1}{n(n-r)} \binom{n}{r+1}^2.$$

Proof. For $k = 2$, Theorem 2.1 yields

$$F_{12}(x, y) = \frac{1}{1 - x + xy - \frac{xy}{1 - x + xy - \frac{xy}{1 - x + xy - \frac{xy}{\dots}}}}.$$

which means that

$$F_{12}(x, y) = \frac{1}{1 - x + xy - xyF_{12}(x, y)}.$$

So the rest it is easy to see. $\square$

Now, we find explicit expressions for $F_{12-3-\dots-k;r}(x)$ in the case $0 \leq r \leq k - 2$.

Following [MV], consider a recurrence relation

$$T_j = \frac{1}{1 - xT_{j-1}}, \quad j \geq 1. \quad (2)$$

The solution of (2) with the initial condition $T_0 = 0$ is denoted by $R_j(x)$, and the solution of (2) with the initial condition

$$T_0 = G_{12-3-\dots-k}(x, y) = \frac{1}{1 - x + xy \binom{k-2}{0} - \frac{xy \binom{k-2}{0}}{1 - x + xy \binom{k-1}{1} - \frac{xy \binom{k-1}{1}}{1 - x + xy \binom{k}{2} - \frac{xy \binom{k}{2}}{\dots}}}$$

is denoted by $S_j(x, y; k)$, or just S_j when the value of k is clear from the context. Our interest in (2) is stipulated by the following relation, which is an easy consequence of Theorem 2.1:

$$F_{12-3-\dots-k}(x, y) = S_{k-2}(x, y; k). \quad (3)$$

Following to [MV, eq. 4], for all $j \geq 1$

$$R_j(x) = \frac{U_{j-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_j\left(\frac{1}{2\sqrt{x}}\right)}, \quad (4)$$

where $U_j(\cos \theta) = \sin(j+1)\theta / \sin \theta$ be the Chebyshev polynomials of the second kind. Next, we find an explicit expression for S_j in terms of G and R_j .

Lemma 2.6. *For any $j \geq 2$ and any $k \geq 2$*

$$S_j(x, y; k) = R_j(x) \frac{1 - xR_{j-1}(x)G_{12-3-\dots-k}(x, y)}{1 - xR_j(x)G_{12-3-\dots-k}(x, y)}. \quad (5)$$

Proof. Indeed, from (2) and $S_0 = G$ we get $S_1 = 1/(1 - xG)$. On the other hand, $R_0 = 0$, $R_1 = 1$, so (5) holds for $j = 1$. Now let $j > 1$, then by induction

$$S_j = \frac{1}{1 - xS_{j-1}} = \frac{1}{1 - xR_{j-1}} \cdot \frac{1 - xR_{j-1}G}{1 - \frac{x(1 - xR_{j-2})R_{j-1}G}{1 - xR_{j-1}}}.$$

Relation (2) for R_j and R_{j-1} yields $(1 - xR_{j-2})R_{j-1} = (1 - xR_{j-1})R_j = 1$, which together with the above formula gives (5). $\square$

As a corollary from Lemma 2.6 and (3) we get the following expression for the generating function $F_{12-3-\dots-k}(x, y)$.

Corollary 2.7. *For any $k \geq 3$*

$$F_{12-3-\dots-k}(x, y) = R_k(x) + (R_{k-2}(x) - R_{k-3}(x)) \sum_{m \geq 1} (xR_{k-2}(x)G_{12-3-\dots-k}(x, y))^m.$$

Now we are ready to express the generating functions $F_{12-3-\dots-k;r}(x)$, $0 \leq r \leq k-2$, via Chebyshev polynomials.

Theorem 2.8. *For any $k \geq 3$, $F_{12-3-\dots-k;r}(x)$ is a rational function given by*

$$\begin{aligned} F_{12-3-\dots-k;r}(x) &= \frac{x^{r-1}U_{k-2}^{r-1}\left(\frac{1}{2\sqrt{x}}\right)}{(1-x)^r U_k^{r+1}\left(\frac{1}{2\sqrt{x}}\right)}, \quad 1 \leq r \leq k-2, \\ F_{12-3-\dots-k;0}(x) &= \frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_k\left(\frac{1}{2\sqrt{x}}\right)}, \end{aligned}$$

where U_j is the j th Chebyshev polynomial of the second kind.

Proof. Observe that $G_{12-3-\dots-k}(x, y) = \frac{1}{1-x} \cdot \frac{1}{1-\frac{x^2}{(1-x)^2}y} + y^{k-1}P(x, y)$, so from Corollary we get

$$\begin{aligned} F_{12-3-\dots-k}(x, y) &= R_k(x) + (R_{k-2}(x) - R_{k-3}(x)) \sum_{m=1}^k \left(\frac{x}{1-x}R_k(x)\right)^m \sum_{n=1}^{k-2} \binom{m-1+n}{n} \frac{x^{2n}}{(1-x)^{2n}} y^n \\ &\quad + y^{k-1}P'(x, y), \end{aligned}$$

where $P(x, y)$ and $P'(x, y)$ are formal power series. To complete the proof, it suffices to use (4) together with the identity $U_{n-1}^2(z) - U_n(z)U_{n-2}(z) = 1$, which follows easily from the trigonometric identity $\sin^2 n\theta - \sin^2 \theta = \sin(n+1)\theta \sin(n-1)\theta$. $\square$

2.2 PATTERN 21-3-...-k

Our second result is a natural analog of the main theorems of [RWZ, MV, Kr].

Theorem 2.9. *For any $k \geq 2$,*

$$F_{21-3-\dots-k}(x, y) = 1 - \frac{x}{xy^{d_1} - \frac{1}{1 - \frac{x}{xy^{d_2} - \frac{1}{1 - \frac{x}{xy^{d_3} - \frac{1}{\ddots}}}}}},$$

where $d_i = \binom{i-1}{k-2}$, and $\binom{a}{b}$ is assumed 0 whenever $a < b$ or $b < 0$.

Proof. Following [MV] we define $\nu_j(\pi)$, $j \geq 3$, as the number of occurrences of 21-3-...-j in π . Define $\nu_2(\pi)$ for any π , as the number of occurrences of 21 in π , $\nu_1(\pi)$ as the number letters of ν , and $\nu_0(\pi) = 1$ for any π , which means that the empty pattern occurs exactly once in each permutation. The *weight* of a permutation π is a monomial in k independent variables $q_1, \dots, q_k$ defined by

$$v_k(\pi) = \prod_{j=1}^k q_j^{\nu_j(\pi)}.$$

The *total weight* is a polynomial

$$V_k(q_1, \dots, q_k) = \sum_{\pi \in \mathcal{S}(1-3-2)} v_k(\pi).$$

The following proposition is implied immediately by the definitions.

Proposition 2.10. $F_{21-3-\dots-k}(x, y) = V_k(x, 1, \dots, 1, y)$ for $k \geq 2$.

We now find a recurrence relation for the numbers $\nu_j(\pi)$. Let $\pi \in S_n$, so that $\pi = (\pi', n, \pi'')$.

Proposition 2.11. For any nonempty $\pi \in \mathcal{S}(1-3-2)$

$$\nu_j(\pi) = \nu_j(\pi') + \nu_j(\pi'') + \nu_{j-1}(\pi'),$$

where $j \neq 2$. Besides, if π'' nonempty then

$$\nu_2(\pi) = \nu_2(\pi') + \nu_2(\pi'') + 1,$$

otherwise

$$\nu_2(\pi) = \nu_2(\pi'').$$

Proof. Similarly to Proposition 2.3 we get π avoids 1-3-2 if and only if π' is a 1-3-2-avoiding permutation of the numbers $\{n-l+1, n-l+2, \dots, n-1\}$, while π'' is a 1-3-2-avoiding permutation of the numbers $\{1, 2, \dots, n-l\}$. Finally, if $(i_1, \dots, i_j)$ is an occurrence of 21-3- $\dots$ - j in π then either $i_j < l$, and so it is also an occurrence of 21-3- $\dots$ - j in π' , or $i_1 > l$, and so it is also an occurrence of 21-3- $\dots$ - j in π'' , or $i_j = l$, and so $(i_1, \dots, i_{j-1})$ is an occurrence of 21-3- $\dots$ -($j-1$) in π' , where $j \neq 2$. For $j = 2$ the proposition is trivial. The result follows. $\square$

Now we are able to find the recurrence relation for the total weight V . Indeed, by Proposition 2.11,

$$\begin{aligned} V_k(q_1, \dots, q_k) &= 1 + \sum_{\emptyset \neq \pi \in \mathcal{S}(1-3-2)} \prod_{j=1}^k q_j^{\nu_j(\pi)} \\ &= 1 + \sum_{\emptyset \neq \pi'' \in \mathcal{S}(1-3-2)} \sum_{\pi' \in \mathcal{S}(1-3-2)} \prod_{j=1}^k q_j^{\nu_j(\pi'')} \cdot q_1^{\nu_1(\pi')+1} q_2 \cdot \prod_{j=2}^{k-1} (q_j q_{j+1})^{\nu_j(\pi')} \cdot q_k^{\nu_k(\pi')} + \\ &\quad + \sum_{\pi' \in \mathcal{S}(1-3-2)} q_1 q_1^{\nu_1(\pi')} q_k^{\nu_k(\pi')} \prod_{j=2}^{k-1} (q_j q_{j+1})^{\nu_j(\pi')}. \end{aligned}$$

Hence

$$\begin{aligned} V_k(q_1, \dots, q_k) &= 1 + q_1 V_k(q_1, q_2 q_3, \dots, q_{k-1} q_k, q_k) + \\ &\quad + q_1 q_2 V_k(q_1, q_2 q_3, \dots, q_{k-1} q_k, q_k) (V_k(q_1, q_2, \dots, q_k) - 1). \end{aligned} \quad (6)$$

Observe now that $V_k(q_1, \dots, q_k) = V_k(q_1, \mathbf{q}^{0,2}, \dots, \mathbf{q}^{0,k})$ and that by (6) and Proposition 2.4

$$\begin{aligned} V_k(q_1, \mathbf{q}^{d,2}, \dots, \mathbf{q}^{d,k}) &= 1 + q_1 V_k(q_1, \mathbf{q}^{d+1,2}, \dots, \mathbf{q}^{d+1,k}) + \\ &\quad + q_1 q^{d,2} V_k(q_1, q^{d+1,2}, \dots, \mathbf{q}^{d+1,k}) (V_k(q_1, \mathbf{q}^{d,2}, \dots, \mathbf{q}^{d,k}) - 1). \end{aligned}$$

To obtain the continued fraction representation for $F_{21-3-\dots-k}(x, y)$ it is enough to use Proposition 2.10 and to observe that

$$q_1 \mathbf{q}^{d,2} \Big|_{q_1=x, q_2=\dots=q_{k-1}=1, q_k=y} = xy^{\binom{d}{k-2}}.$$

$\square$

Corollary 2.12.

$$F_{21}(x, y) = \frac{1 - x + xy - \sqrt{(1-x)^2 - 2x(1+x)y + x^2y^2}}{2xy},$$

in other words, for any $r \geq 1$

$$f_{21;r}(n) = \frac{r+1}{n(n-r)} \binom{n}{r+1}^2.$$

Proof. For $k = 2$, $q_1 = x$ and $q_2 = y$, Proposition 2.10 and (6) yields $F_{21}(x, y) = 1 + xF_{21}(x, y) + xyF_{21}(x, y)(F_{21}(x, y) - 1)$, which means $F_{21}(x, y) = F_{12}(x, y)$. The rest of the proof obtained by Corollary 2.5. $\square$

Now, we are ready to find an explicit expressions for $F_{21-3-\dots-k;r}(x)$ in the case $0 \leq r \leq k-2$.

Consider a recurrence relation

$$T'_j = 1 - \frac{x}{x - \frac{1}{T'_{j-1}(x)}}, \quad j \geq 1. \quad (7)$$

The solution of (7) with the initial condition $T'_0 = 0$ is given by $R_j(x)$ (Lemma 2.13), and the solution of (7) with the initial condition

$$T'_0 = G_{21-3-\dots-k}(x, y) = \frac{1}{xy^{\binom{k-2}{k-2}} - \frac{1}{1 - \frac{1}{xy^{\binom{k-1}{k-2}} - \frac{1}{1 - \frac{1}{xy^{\binom{k}{k-2}} - \frac{1}{\ddots}}}}}},$$

is denoted by $S'_j(x, y; k)$, or just S'_j when the value of k is clear from the context. Our interest in (7) is stipulated by the following relation, which is an easy consequence of Theorem 2.9:

$$F_{21-3-\dots-k}(x, y) = S'_k(x, y; k). \quad (8)$$

First of all, we find an explicit formula for the functions $T'_j(x)$ in (7).

Lemma 2.13. For any $j \geq 1$,

$$T'_j(x) = R_j(x). \quad (9)$$

Proof. Indeed, it follows immediately from (7) that $T'_0(x) = 0$ and $T'_1(x) = 1$. Let us induction, we assume $T'_{j-1}(x) = R_{j-1}(x)$, and prove that $T'_j(x) = R_j(x)$. By use (7)

$$T'_j(x) = 1 - \frac{x}{x - \frac{1}{R_{j-1}(x)}}.$$

On the other hand, following to [MV], $R_j(x) = \frac{1}{1 - xR_{j-1}(x)}$ which means that $R_j(x) = 1 + xR_{j-1}(x)R_j(x)$, hence $T'_j(x) = R_j(x)$. $\square$

Next, we find an explicit expression for S'_j in terms of G and R_j .

Lemma 2.14. *For any $j \geq 2$ and any $k \geq 2$*

$$S'_j(x, y; k) = R_j(x) \frac{1 - xR_{j-1}(x)G_{21-3-\dots-k}(x, y; k)}{1 - xR_j(x)G_{21-3-\dots-k}(x, y)}. \quad (10)$$

As a corollary from Lemma 2.14 and (6) we get the following expression for the generating function $F_{21-3-\dots-k}(x, y)$.

Corollary 2.15. *For any $k \geq 3$*

$$F_{21-3-\dots-k}(x, y) = R_k(x) + (R_{k-2}(x) - R_{k-3}(x)) \sum_{m \geq 1} (xR_{k-2}(x)G_{21-3-\dots-k}(x, y))^m.$$

Now we are ready to express the generating functions $F_{21-3-\dots-k;r}(x)$, $0 \leq r \leq k-2$, via Chebyshev polynomials.

Theorem 2.16. *For any $k \geq 3$, $F_{21-3-\dots-k;r}(x)$ is a rational function given by*

$$\begin{aligned} F_{21-3-\dots-k;r}(x) &= \frac{x^{\frac{r-1}{2}} U_{k-2}^{r-1}\left(\frac{1}{2\sqrt{x}}\right)}{U_{k-1}^{r+1}\left(\frac{1}{2\sqrt{x}}\right)}, \quad 1 \leq r \leq k-2, \\ F_{21-3-\dots-k;0}(x) &= \frac{U_{k-1}\left(\frac{1}{2\sqrt{x}}\right)}{\sqrt{x}U_k\left(\frac{1}{2\sqrt{x}}\right)}, \end{aligned}$$

where U_j is the j th Chebyshev polynomial of the second kind.

Proof. Observe that $G_{21-3-\dots-k}(x, y) = 1 + \frac{x}{1-x-xy} + y^{k-1}P(x, y)$, so from Corollary 2.15 we get

$$F_{21-3-\dots-k}(x, y) = R_k(x) + (R_{k-2}(x) - R_{k-3}(x)) \sum_{m=1}^k \left(xR_{k-2}(x)\left(1 + \frac{x}{1-x-xy}\right)\right)^m + y^{k-1}P'(x, y),$$

where $P(x, y)$ and $P'(x, y)$ are formal power series. To complete the proof, it suffices to use (9) together with the identity $U_{n-1}^2(z) - U_n(z)U_{n-2}(z) = 1$. $\square$

Remark 2.17. *Theorem 2.16 and [MV] yields the number of 1-3-2-avoiding permutations in S_n such that containing r times the pattern $21-3-\dots-k$ is the same number of 1-3-2-avoiding permutations in S_n such that containing r times the pattern $1-2-3-\dots-k$, for all $r = 0, 1, 2, \dots, k-2$. However, the question if there exist a natural bijection between the set of 1-3-2-avoiding permutations in S_n such containing r times the generalized pattern $21-3-\dots-k$, and the set of 1-3-2-avoiding permutations in S_n such containing r times the classically pattern $1-2-3-\dots-k$.*

2.3 PATTERNS: $\tau = 12\dots k$ AND $\tau = k\dots 21$

Let $\pi \in S_n$; we say π has d -increasing canonical decomposition if π has the following form

$$\pi = (\pi^1, \pi^2, \dots, \pi^d, a_d, \dots, a_2, a_1, n, \pi^{d+1}),$$

where all the entries of π^i are greater than all the entries of π^{i+1} , and $a_d < a_{d-1} < \dots < a_1 < n$. We say π has d -decreasing canonical decomposition if π has the following form

$$\pi = (\pi^1, n, a_1, \dots, a_d, \pi^{d+1}, \pi^d, \dots, \pi^1),$$

where all the entries of π^i are greater than all the entries of π^{i+1} , and $a_d < a_{d-1} < \dots < a_1 < n$. The following proposition is the base of all the other results in this Section.

Proposition 2.18. *Let $\pi \in S_n(1-3-2)$. Then there exists unique $d \geq 0$ and $e \geq 0$ such that π has a d -increasing canonical decomposition, and has e -decreasing canonical decomposition.*

Proof. Let $\pi \in S_n(1-3-2)$, and let $a_d, a_{d-1}, \dots, a_1, n$ a maximal increasing subsequence of π such that $\pi = (\pi', a_d, \dots, a_1, n, \pi'')$. Since π avoids 1-3-2 there exists d subsequences π^j such that $\pi = (\pi^1, \dots, \pi^d, a_d, \dots, a_1, n, \pi'')$, and all the entries of π^i are greater than all the entries of π^{i+1} , and all the entries of π^d greater than all entries of π'' . Hence, π has d -increasing canonical decomposition. Similarly, there exist e unique such that π is e -decreasing canonical decomposition. $\square$

Let us define $I_\tau(x, y; d)$ (respectively; $J_\tau(x, y; e)$) be the generating function for all d -increasing (respectively; e -decreasing) canonical decomposition of permutations in $S_n(1-3-2)$ with exactly r occurrences of τ . The following proposition is implied immediately by the definitions.

Proposition 2.19.

$$F_\tau(x, y) = 1 + \sum_{d \geq 0} I_\tau(x, y; d) = 1 + \sum_{e \geq 0} J_\tau(x, y; e).$$

Proof. Immetaitley, by definitions of the generating functions and Proposition 2.18 (1 for the empty permutation). $\square$

Now, we present an examples for Proposition 2.18 and Proposition 2.19.

First example: $F_{12\dots k}(x)$ and $F_{k\dots 21}(x)$.

Theorem 2.20. $F_{k\dots 21}(x, y) = F_{12\dots k}(x, y)$, such that

$$F_{12\dots k}(x, y) = \sum_{n=0}^{k-2} x^n F_{12\dots k}^n(x, y) + \frac{x^{k-1} F_{12\dots k}^{k-1}(x, y)}{1 - xy F_{12\dots k}(x, y)}.$$

Proof. By Proposition 2.18 and definitions it is easy to obtain for all $d \geq 0$

$$I_{12\dots k}(x, y; d) = x^{d+1} y^{s_d} F_{12\dots k}^{d+1}(x, y),$$

where $s_d = d + 1 - k$ for $d \geq k - 1$, and otherwise $s_d = 0$. So by Proposition 2.19 the theorem holds.

Similarly, we obtain the same result for $F_{k\dots 21}(x, y)$. $\square$

As a remark, by the above theorem, it is easy to obtain the same Corollaries 2.5 and 2.12 results.

Second example. Here we calculate $F_{1-2-\dots-(l-1)-l(l+1)\dots k}(x, y)$ where $l \leq k - 1$.

Theorem 2.21. *Let $1 \leq l \leq k - 1$. Then $F_{1-2-\dots-(l-1)-l(l+1)\dots k}(x, y) = U_l(x, 1, \dots, 1, y)$ where*

$$U_l(q_1, \dots, q_l) = 1 + \sum_{d \geq 0} \left((q_l^{\binom{d+1+l-k}{l}} \prod_{j=1}^{l-1} q_j^{\binom{d+1}{j}} \prod_{j=0}^d U_l(p_{1;j}, \dots, p_{l;j}) \right),$$

and for $i = 1, 2, \dots, l$, $p_{i;j} = \prod_{m=1}^{l-1} q_j^{\binom{j}{m-i}}$, $p_{l;j} = q_l$ for all $0 \leq j \leq k - l$, and $p_{i;j} = \prod_{m=1}^l p_{i;k-l}^{\binom{j-k+l}{l-i}}$ for all $j \geq k - l + 1$.

Proof. Following [MV] we define $\gamma_j(\pi)$, $j \leq l-1$, as the number of occurrences of $1-2-\dots-j$ in π . Define $\gamma_l(\pi)$ for any π , as the number of occurrences of $1-2-\dots-(l-1)-l(l+1)\dots k$ in π , and $\gamma_0(\pi) = 1$ for any π , which means that the empty pattern occurs exactly once in each permutation. The *weight* of a permutation π is a monomial in l independent variables $q_1, \dots, q_l$ defined by

$$u_l(\pi) = \prod_{j=1}^l q_j^{\gamma_j(\pi)}.$$

The *total weight* is a polynomial

$$U_l(q_1, \dots, q_l) = \sum_{\pi \in \mathcal{S}(1-3-2)} u_l(\pi).$$

The following proposition is implied immediately by the definitions, and Proposition 2.18.

Proposition 2.22. $F_{1-2-\dots-(l-1)-l(l+1)\dots k}(x, y) = U_k(x, 1, \dots, 1, y)$ for $k > l \geq 1$, and $U_l(q_1, \dots, q_l) = 1 + \sum_{d \geq 0} \sum_{\pi \in A_d} u_l(\pi)$, where A_d is the set of all d -increasing canonical decomposition permutations in $\mathcal{S}(1-3-2)$.

Let us denote $U_{l;d}(q_1, \dots, q_l) = \sum_{\pi \in A_d} u_l(\pi)$.

Proposition 2.23. For any $d \geq 0$,

$$U_{l;d}(q_1, \dots, q_l) = q_l^{\binom{d+1+l-k}{l}} \prod_{j=1}^{l-1} q_j^{\binom{d+1}{j}} \prod_{j=0}^d U_l(p_{1;j}, \dots, p_{l;j}).$$

Proof. Let π is d -increasing canonical decomposition; that is,

$$\pi = (\pi^1, \pi^2, \dots, \pi^d, a_d, \dots, a_2, a_1, n, \pi^{d+1}),$$

where the numbers $a_d < a_{d-1} < \dots < a_1 < n$ are appear as consecutive numbers in π , the all entries of π^j are greater than all the entries of π^{j+1} , and the all entries of π^d greater than a_d . So, by calculate $u_l(\pi)$ and sum over all $\pi \in A_d$ we have that

$$U_{l;d}(q_1, \dots, q_d) = q_l^{\binom{d+1+l-k}{l}} \cdot \prod_{j=1}^{l-1} q_j^{\binom{d+1}{j}} \cdot \prod_{j=0}^d U_l(p_{1;j}, \dots, p_{l;j}).$$

□

Therefore, Theorem 2.21 holds, by use the above results; Proposition 2.22 and Proposition 2.23. □

Now, let $l = k - 1$ and by use Theorem 2.21, it is easy to obtain the following.

Corollary 2.24. For $k \geq 3$,

$$F_{1-2-\dots-(k-2)-(k-1)k}(x, y) = \sum_{j=0}^{k-1} (x F_{1-2-\dots-(k-2)-(k-1)k}(x, y))^j.$$

Remark 2.25. Similarly, the argument of d -increasing canonical decomposition, or the argumnet d -decreasing canonical decomposition yields others formulas, for example, formula for $F_{12-3-45}(x, y)$.

3. THREE LETTERS PATTERN WITHOUT INTERNAL DASHES

In this section, we give a complete answer for $F_\tau(x, y)$ where τ is a generalized pattern without internal dashes; that is, τ is 123, 213, 231, 312, and 321. This by the following four subsections.

3.1 Patterns 123 and 321.

Theorem 3.1.

$$F_{123}(x, y) = F_{321}(x, y) = \frac{1 + xy - x + \sqrt{1 - 2x - 3x^2 - xy(2 - 2x - xy)}}{2x(x + y - xy)}.$$

Proof. Theorem 2.20 yields, $F_{123}(x, y) = F_{321}(x, y) = H$ where

$$H = 1 + xH + \frac{x^2 H^2}{1 - xyH},$$

and the rest it is easy to see. □

3.2 Pattern 231.

Theorem 3.2.

$$F_{231}(x, y) = \frac{1 - 2x + 2xy - \sqrt{1 - 4x + 4x^2 - 4x^2 y}}{2xy},$$

that is, for all $r, n \geq 0$

$$F_{231;r}(x) = \frac{1}{r+1} \binom{2r}{r} \frac{x^{2r+1}}{(1-2x)^{2r+1}}, \quad f_{231;r}(n) = \frac{2^{n-2r-1}}{r+1} \binom{n-1}{2r} \binom{2r}{r}.$$

Proof. Let $l = \pi^{-1}(n)$. Since π avoids 1-3-2, each number in π' is greater than any of the numbers in π'' . Therefore, π' is a 1-3-2-avoiding permutation of the numbers $\{n-l+1, n-l+2, \dots, n-1\}$, while π'' is a 1-3-2-avoiding permutation of the numbers $\{1, 2, \dots, n-l\}$. On the other hand, if π' is an arbitrary 1-3-2-avoiding permutation of the numbers $\{n-l+1, n-l+2, \dots, n-1\}$ and π'' is an arbitrary 1-3-2-avoiding permutation of the numbers $\{1, 2, \dots, n-l\}$, then $\pi = (\pi', n, \pi'')$ is 1-3-2-avoiding.

Now let us observe all possibilities of π' and π'' are empty or not. This yields

$$F_{231}(x, y) = 1 + x + 2x(F_{231}(x, y) - 1) + xy(F_{231}(x, y) - 1)^2,$$

hence, the rest it is easy to see. □

3.3 Pattern 213.

Theorem 3.3.

$$F_{213}(x, y) = \frac{1 - x^2 + x^2 y - \sqrt{1 + 2x^2 - 2x^2 y + x^4 - 2x^4 y + x^4 y^2 - 4x}}{2x(1 + xy - x)}.$$

Proof. Let $D(x, y)$ be the generating function of all 1-3-2-avoiding permutations $(\alpha', n) \in S_n$ such that containing 213 exactly r times. Let $\alpha = (\alpha', n, \alpha'')$; if we consider the two cases α' empty or not we obtain $F_{213}(x, y) = 1 + D(x, y)F_{213}(x, y)$. Now let $\alpha = (\alpha', n)$; if we observe the two cases α' empty or not, then similarly it is easy to see

$$D(x, y) = x + x^2 + x^2 y(F_{213}(x, y) - 1) + x^2(D(x, y) - 1) + x^2(D(x, y) - 1)(F_{213}(x, y) - 1).$$

However,

$$F_{213}(x, y) = 1 + xF_{213}(x, y) \frac{1 + x - xy + x(y-1)F_{213}(x, y)}{1 - xF_{213}(x, y)},$$

hence, the rest it is easy to see. $\square$

3.4 Pattern 312.

Theorem 3.4.

$$F_{312}(x, y) = \frac{1 - x^2 + x^2y - \sqrt{1 + 2x^2 - 2x^2y + x^4 - 2x^4y + x^4y^2 - 4x}}{2x(1 + xy - x)}.$$

Proof. Let $\alpha \in \mathcal{S}(1-3-2)$; if $\alpha = \emptyset$, then there one permutation, otherwise by Proposition 2.18 we can write $\alpha = (\alpha^1, n, a_1, a_2, \dots, a_d, \alpha^{d+1}, \alpha^d, \dots, \alpha^2)$ where all the entries of α^j greater than all the entries of α^{j+1} , and $n > a_1 > a_2 > \dots > a_d$. Hence, for any $d = 0, 1$ the generating function of these permutations in these cases is $x^{d+1}F_{312}(x, y)$. Let $d \geq 2$, If $\alpha^{d+1} = \emptyset$, then the generating function of these permutations in this case is $x^{d+1}F_{312}^d(x, y)$, otherwise the generating function is $x^{d+1}yF_{312}^d(x, y)(F_{312}(x, y) - 1)$. Hence

$$F_{312}(x, y) = 1 + (x + x^2)F_{312}(x, y) + \sum_{d \geq 2} x^{d+1}F_{312}^d(x, y) + \sum_{d \geq 2} x^{d+1}yF_{312}^d(x, y)(F_{312}(x, y) - 1),$$

which means that

$$F_{312}(x, y) = 1 + xF_{312}(x, y) + \frac{x^2F_{312}(x, y)}{1 - xF_{312}(x, y)} + \frac{x^2yF_{312}(x, y)(F_{312}(x, y) - 1)}{1 - xF_{312}(x, y)},$$

so the rest it is easy to see. $\square$

4. SOME OF 3-LETTER PATTERNS WITH ONE DASH

In this section, we present some of examples $F_\tau(x, y)$ where τ is a generalized pattern with one dash. Theorem 2.1 yields as follows.

Theorem 4.1. *The generating function $F_{12-3}(x, y)$ is given by the continued fraction*

$$\cfrac{1}{1 - \cfrac{x}{1 - x + xy - \cfrac{xy}{1 - x + xy^2 - \cfrac{xy^2}{\ddots}}}}.$$

Theorem 2.9 yields as follows.

Theorem 4.2. *For any $k \geq 2$,*

$$F_{21-3}(x, y) = 1 - \cfrac{x}{x - \cfrac{1}{1 - \cfrac{x}{xy - \cfrac{1}{1 - \cfrac{x}{xy^2 - \cfrac{1}{\ddots}}}}}}.$$

For $k = 3$ and $l = 2$ Theorem 2.24 yields the following.

Theorem 4.3.

$$F_{1-23}(x, y) = 1 + xF_{1-23}(x, y) + \sum_{d \geq 1} x^{d+1} y^{\binom{d}{2}} F_{1-23}(x, y) \prod_{j=0}^{d-1} F_{1-23}(xy^j, y).$$

Corollary 4.4.

$$\begin{aligned} F_{1-23;0}(x) &= \frac{1-x-\sqrt{1-2x-3x^2}}{2x^2}; \\ F_{1-23;1}(x) &= \frac{x-1}{2x} + \frac{1-2x-x^2}{2x\sqrt{1-2x-3x^2}}; \\ F_{1-23;2}(x) &= \frac{x^4}{(1-2x-3x^2)^{3/2}}; \\ F_{1-23;3}(x) &= x^2 - 1 + \frac{11x^7+43x^6+41x^5-7x^4-25x^3+x^2+5x-1}{(1-2x-3x^2)^{5/2}}. \end{aligned}$$

Proof. By Theorem 4.3 with use $F_{1-23}(x, 0) = F_{1-23;0}(x)$ we get

$$F_{1-23;0}(x) = 1 + xF_{1-23;0}(x) + x^2 F_{1-23;0}^2(x),$$

which means the first formula holds.

By Theorem 4.3 we get

$$\frac{d}{dy} F_{1-23}(x, 0) = x \frac{d}{dy} F_{1-23}(x, 0) + 2x^2 F_{1-23}(x, 0) \frac{d}{dy} F_{1-23}(x, 0) + x^3 F_{1-23}(x, 0)^2 F_{1-23}(0, 0),$$

and with use $F_{1-23;1}(x) = \frac{d}{dy} F_{1-23}(x, y)|_{y=0}$ and the first formula, the second formula holds.

Similarly, by Theorem 4.3 and use $F_{1-23;r}(x) = \frac{1}{r!} \frac{d^r}{dy^r} F_{1-23}(x, y)|_{y=0}$ the others formulas holds. $\square$

Theorem 4.5.

$$F_{2-13}(x, y) = \frac{1}{1 - \frac{x}{1 - \frac{xy}{1 - \frac{xy}{1 - \frac{xy^2}{1 - \frac{xy^2}{1 - \dots}}}}}}.$$

Proof. By use Proposition 2.18 and Proposition 2.19, we obtain

$$F_{2-13}(x, y) = 1 + xF_{2-13}(x, y) \sum_{d \geq 0} x^d F_{2-13}^d(xy, y),$$

and the rest is easy to see. $\square$

5. FRUTHER RESULTS

First of all, let us denote by $G_{\tau;\phi}(x, y)$ the generating function for the number of permutations in $S_n(1-3-2, \tau)$ such containing ϕ exactly r times; that is

$$G_{\tau;\phi}(x, z) = \sum_{n \geq 0} x^n \sum_{\pi \in S_n(1-3-2, \tau)} y^{a_\phi(\pi)},$$

where $a_\phi(\pi)$ is the number of occurrences of ϕ in π . In this section, (similarly to above sections) we find $G_{\tau;\phi}(x, y)$ in terms of continued fractions or by explicit formula, for some cases of τ and ϕ .

Theorem 5.1. *The generating functions $G_{123;213}(x, y)$ and $G_{321;312}(x, y)$ are given by*

$$\frac{1}{1 - x - x^2(1 - y) - \frac{x^2y}{1 - x - x^2(1 - y) - \frac{x^2y}{1 - x - x^2(1 - y) - \frac{x^2y}{\ddots}}}},$$

or,

$$\frac{1 - x - x^2 + x^2y - \sqrt{(1 - x - x^2)^2 - 2yx^2(1 + x + x^2) + x^4y^2}}{2x^2y}.$$

Theorem 5.2.

$$G_{123;231}(x, y) = H(x, y) + x^2(1 - y)H(x, y)^2,$$

where $H(x, y) = \frac{1}{1 - x - x^2yH(x, y)}$, which means number of permutations in $S_n(1\text{-}3\text{-}2, 123)$ such contains 231 exactly $r \geq 1$ times is given by

$$C_r \left(\binom{n}{2r} + \binom{n-1}{2r+1} \right) + C_{r-1} \binom{n-2}{2r-1},$$

where C_r is the r th Catalan number, and for $r = 0$ is given by n .

Theorem 5.3. *The generating functions $G_{213;123}(x, y)$ and $G_{312;321}(x, y)$ given by*

$$\frac{1 - x - x^2 + xy - \sqrt{(1 - x - x^2)^2 - 2xy(1 - x + x^2) + x^2y^2}}{2xy(1 - x)}.$$

As a concluding remark we note that are many questions left to answer. If there exist a bijection between, for example, the set of 1-3-2-avoiding permutations in S_n such containing r times the generalized pattern $21\text{-}3\text{-}\dots\text{-}k$, and the set of 1-3-2-avoiding permutations in S_n such that containing r times the classically pattern $1\text{-}2\text{-}3\text{-}\dots\text{-}k$, where $r = 0, 1, \dots, k - 2$.

REFERENCES

- [BS] E. BABSON AND E. STEINGRIMSSON, Generalized permutation patterns and a classification of the Mahonian statistics, *Séminaire Lotharingien de Combinatoire*, B44b:18pp, (2000).
- [C] A. CLAEISSON, Generalised pattern avoidance, *European Journal of Combinatorics*, **22** (2001) 961–973.
- [CM] A. CLAEISSON AND T. MANSOUR, Permutations avoiding a pair of generalized patterns of length three with exactly one dash, preprint CO/0107044.
- [CW] T. CHOW AND J. WEST, Forbidden subsequences and Chebyshev polynomials, *Discr. Math.*, **204** (1999) 119–128.
- [EN] S. ELIZALDE AND M. NOY, Enumeration of subwords in permutations. In *Formal power series and algebraic combinatorics (Tempe, 2001)*, Arizona State University 2001, 179–189.
- [JR] M. JANI AND R.G. RIEPER, Continued fractions and catalan problems, *Elect. J. Combin.* **7** (2000) #R1.
- [Ki] S. KITAEV, Multi-Avoidance of generalised patterns, to appear.
- [Kn] D. KNUTH, *The Art of Computer Programming*, volume 3, Addison Wesley, Reading, MA 1973.

- [Kr] C. KRATTENTHALER, Permutations with restricted patterns and Dyck paths, (2000), preprint CO/0002200.
- [LS] V. LAKSHMIBAI AND B. SANDHYA, Criterion for smoothness of Schubert varieties in $Sl(n)/B$, *Proc. Indian Acad. Sci.* **100** no. 1 (1990) 45–52.
- [LW] L. LORENTZEN AND H. WAADELAND, Continued fractions with applications, North-Holland 1992.
- [MV] T. MANSOUR AND A. VAINSHTEIN, Restricted permutations, continued fractions, and Chebyshev polynomials, *Electron. J. Combin.*, **7** (2000) #R17.
- [N] J. NOONAN, The number of permutations containing exactly one increasing subsequence of length three, *Discr. Math.* **152** (1996) 307–313.
- [NZ] J. NOONAN AND D. ZEILBERGER, The enumeration of permutations with a prescribed number of “forbidden” patterns, *Adv. Appl. Math.* **17** (1996) 381–407.
- [SS] R. SIMION AND F. SCHMIDT, Restricted permutations *Europ. J. Comb.* **6** (1985) 383–406.
- [RWZ] A. ROBERTSON, H. WILF, AND D. ZEILBERGER, Permutation patterns and continuous fractions, *Elec. J. Comb.* **6** no. 1 (1999) #R38.
- [W] J. WEST, Generating trees and the Catalan and Schröder numbers, *Discr. Math.* **146** (1995) 247–262.